

FUNGOUT®

Fungicide & Bactericide

*For use on outdoor food,
nonfood and greenhouse*

Disease Prevention for several crops



0 Day PHI

4 Hours REI



TECHNICAL SPECIFICATIONS

- Active ingredients: Citric acid
- Packaging: One (1) case = 2 x 2.5 gal
- Formulation Type: Liquid concentrate
- Chemical Group: N/A Biopesticide
- PHI: 0 days
- REI: When spray is dry or 4 hours

MODE OF ACTION

FungOUT forms a protective barrier on the surface of the leaves that inhibit pathogen development. It also has antifungal effects on conidia and ascospores germination, germ tube formation and elongation, appressorium formation, hyphae branching and haustorium formation. In addition, it has a direct bactericide activity and also reduces bacterial growth and the appearance of various disease symptoms.

Manufactured by:

925 des Calfats • Lévis (QC) G6Y 9E8

Toll free: 1 (866) 622-3222 • Tel.: (418) 838-4441 • Fax: (418) 838-9909

info@aefglobal.com • www.aefglobal.com

Update : 26/09/2018



CROP	DISEASE	RATE (% v/v)
Crop groups 11 and 12 Pome and Stone Fruits such as: Apples, Pears, Quince, Prunes, Apricots, Cherries, Nectarines, Peaches, Plums	Powdery mildew (<i>Podosphaera leucotricha</i> , <i>Podosphaera clandestina</i>) Apple scab (<i>Venturia inaequalis</i>)	1.2% v/v
	Fire blight (<i>Erwinia amylovora</i>)	1.4% v/v (with a non-ionic surfactant)
Grapes	Powdery mildew (<i>Erysiphe necator</i>)	2.1% v/v
Field Cucurbits Crop Group 9 (such as: melon, squashes, pumpkins and cucumbers)	Powdery mildew (<i>Podosphaera fusca</i> ; <i>Erysiphe cichoracearum</i> , syn. <i>Golovinomyces cichoracearum</i>)	2.1% v/v
Field Tomatoes	Bacterial canker (<i>Clavibacter michiganensis</i>)	2.1% v/v
	Powdery mildew (<i>Oidium neolycopersici</i>)	1.2 to 1.6% v/v
Spinach (Organic)	Downy mildew (<i>Peronospora farinosa</i>)	2.1% v/v
Hops	Powdery mildew (<i>Podosphaera macularis</i>)	1.2% v/v
	Downy mildew (<i>Pseudoperonospora humuli</i>)	2.1% v/v
Citrus Fruits Crop group 10-10 such as: Oranges, Lemons, Limes, Grapefruit	Bacterial canker (<i>Xanthomonas spp.</i>)	2.1% v/v
Tobacco	Target spot <i>Rhizoctonia solani</i>	2.1% v/v
Greenhouse Tomatoes	Bacterial canker (<i>Clavibacter michiganensis</i>)	2.1% v/v
	Powdery mildew (<i>Oidium neolycopersici</i>)	1.2 to 1.6% v/v
Greenhouse Cucurbits Crop Group 9 (such as cucumbers)	Powdery mildew (<i>Podosphaera fusca</i> ; <i>Erysiphe cichoracearum</i> , syn. <i>Golovinomyces cichoracearum</i>)	2.1% v/v
Greenhouse and outdoor ornamentals (including cut flowers)	Powdery mildew (<i>Podosphaera fusca</i> ; <i>Erysiphe cichoracearum</i> , <i>oidium sp.</i> <i>Sawadaea sp.</i>)	1.2 to 1.6% v/v
Greenhouse and outdoor roses	Black spot (<i>Diplocarpon rosae</i>) Powdery mildew (<i>Podosphaera pannosa</i>)	1.2 to 1.6% v/v
Greenhouse and field leafy vegetables Crop group 4 such as: Celery, head lettuce and leaf lettuce	Powdery mildew (<i>Erysiphe spp.</i> , <i>Podosphaera fuliginea</i> , syn. <i>Sphaerotheca fuliginea</i>)	1.2 to 1.6% v/v
Indoor and outdoor medical and recreational cannabis (marijuana)	Powdery mildew (<i>Sphaerotheca macularis</i>)	1.2 to 1.6% v/v

FungOUT has not been registered by the United States Environmental Protection Agency. AEF Global Inc. represents that this product qualifies for exemption from registration under the Federal Insecticide, Fungicide and Rodenticide Act. Please refer to product label for complete application details.

Always read and follow label directions. Log on to www.aefglobal.com for full label information.

Manufactured by:
 925 des Calfats • Lévis (QC) G6Y 9E8
 Toll free: 1 (866) 622-3222 • Tel.: (418) 838-4441 • Fax: (418) 838-9909
info@aefglobal.com • www.aefglobal.com