

Bora WP

*For the Suppression of Root Diseases caused by Pythium, Rhizoctonia and Fusarium in
Greenhouse Crops and Outdoor Nursery Plants*

COMMERCIAL

WETTABLE POWDER

READ THE LABEL BEFORE USING

GUARANTEE: *Trichoderma harzianum* Rifai strain KRL-AG2 contains a minimum of 1.0×10^7 colony forming units per gram of dry weight

Warning, contains the allergens wheat and sulfites.

POTENTIAL SENSITIZER

LIVE ORGANISMS

REGISTRATION No. 31104 PEST CONTROL PRODUCTS ACT

NET CONTENTS: 0.45 – 90.9 kg

AEF GLOBAL Inc. 925 des Calfats, Lévis, QC G6Y 9E8

(418) 838-4441

Toll free help-line: 1-866-622-3222

www.aefglobal.com

Lot No.:

Date of Manufacture: Use within 12 months of date of manufacture.

GENERAL INFORMATION: Bora WP is a biological fungicide for the suppression of root diseases in greenhouse crops and outdoor nursery plants. The active ingredient is a beneficial microbe *Trichoderma harzianum* strain T-22 (KRL-AG2). When applied to transplants, greenhouse planting mix or soil, Bora WP actively grows onto plant roots as they develop and protects roots against *Pythium*, *Rhizoctonia* and *Fusarium*. Bora WP can be used alone or in conjunction with certain chemical fungicides; contact manufacturer or distributor for compatibility Technical Bulletin.

For control of seed rot and damping off of internally infected cuttings, use chemical fungicides or other appropriate measures in addition to Bora WP.

Note: Bora WP contains live spores of a beneficial microbe. Bora WP becomes active when soil temperatures are above 10 degrees C. Control is not effective while soils remain cold and is more effective in neutral or acidic soils than in alkaline soils.

DIRECTIONS FOR USE:

DRENCH APPLICATION TO POTTING MIX OR SOIL

Greenhouse Peppers, Tomatoes, and Cucumbers
Greenhouse Ornamental Crops
Outdoor Nursery Crops

POTTING MIX: Suspend Bora WP in sufficient water (eg. 30-45 g/100L) to achieve uniform application and apply at the rate of 55-110 g per cubic metre (loose) of greenhouse potting mix, soil or planting beds. Bora WP can be applied through low pressure watering nozzles such as fan nozzles or other watering systems. Agitate to maintain suspension.

NURSERY DRENCH:

Suspend Bora WP in sufficient water (eg. 30 – 45 g/100 L) to achieve uniform application and apply at the rate of 55 – 110 g per cubic metre (loose) of nursery potting mix, soil or planting beds. Bora WP can be applied through low pressure watering nozzles such as fan nozzles or other watering systems. Agitate to maintain suspension.

DIP APPLICATION

Bulbs: All Ornamental Crops

Dip bulbs in suspension of 60 g/L Bora WP prior to planting.

It is not possible to test all ornamental varieties, and some may show delayed growth. Bora WP should first be tested on small lot of plants, if the response of the variety is not known, before using it for commercial scale production.

GENERAL APPLICATION INSTRUCTIONS

Bora WP can be applied using hand-held backpack or ground spray equipment. **AGITATION IS REQUIRED TO MAINTAIN SUSPENSION.** Application equipment should be cleaned before use of this product, and prepared sprays should be used within 4 hours of preparation. To prevent nozzle tip clogging the manufacturer recommends using 50 – 100 mesh inlet screens in all spray equipment.

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN.

May cause sensitization. Avoid contact with skin, eyes or clothing. Avoid breathing dust or spray mist. Workers and handlers (includes mixer/loader and applicators) must wear a long sleeved shirt, long pants, shoes plus socks, waterproof gloves and eye goggles as well as a dust/mist filtering respirator (MSHA/NIOSH approval number prefix TC-21C) or a NIOSH approved respirator with any N-95, R-95, P-95 or HE filter for biological products when mixing/loading or applying the product and until mists have settled and during all clean-up/repair activities. Wash thoroughly with soap and water after handling product. Remove contaminated clothing and follow manufacturer's directions for cleaning/maintaining personnel protective equipment (PPE) before use. If no such instructions are available use clothing detergent and hot water for cleaning all washable PPE. Keep and wash PPE separately from other laundry. Wash hands after use

ENVIRONMENTAL HAZARDS:

As this pesticide is not registered for the control of pests in aquatic systems, DO NOT use to control aquatic pests.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

To reduce runoff from treated areas into aquatic habitats, consider the characteristics and conditions of the site before treatment. Site characteristics and conditions that may lead to runoff include, but are not limited to: heavy rainfall, moderate to steep slope, bare soil, poorly draining soil (e.g. soils that are compacted, fine textured, or low in organic matter such as clay).

DO NOT apply by air.

FIRST AID:

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

GENERAL: Seek medical attention immediately if irritation or signs of toxicity occur and persist or is severe. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION: Treat symptomatically.

STORAGE:

Store in a secure dry place at temperatures between 2°C and 5°C. Do not contaminate water, food, or feed during storage. Keep product in original container during storage and keep container lid tightly closed when not in use. This product should be used within 12 months of the date of manufacture when stored under refrigerated conditions.

DISPOSAL/ DECONTAMINATION:

1. Thoroughly empty the contents of the container into the application device or medium.
2. Make the empty container unsuitable for further use.
3. Dispose of the container in accordance with provincial requirements.
4. For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER:

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use this product.