



Buran

Fongicide

Buran is a fungicide for use in Orchards and Vineyards. Include Buran in your disease management program for scab and powdery mildew.

Disease Prevention for Orchards and Vineyards

FEATURES AND BENEFITS

- ✓ **Zero** days to harvest, allows to keep control of diseases and maintain quality
- ✓ **Safe** for use with beneficial and pollinators when used according to label instructions
- ✓ **Minimum risk** of resistance
- ✓ **Perfect integration** in your IPM program
- ✓ **Great alternative** to use in rotation with other products or in tank mixes
- ✓ **Approved** for organic agriculture

TECHNICAL SPECIFICATIONS

- **Canadian registration number:** 30601
- **Active ingredients:** Garlic
- **Packaging:** One (1) case = 2 x 10L
- **Formulation Type:** Liquid concentrate
- **Chemical Group:** N/A Biopesticide
- **PHI:** 0 day
- **REI:** When spray is dry

RECOMMANDATIONS POUR DES RÉSULTATS OPTIMAUX

- Use minimum spray volume to obtain a complete coverage of foliage, flowers and fruits
- Apply every 7-10 days when conditions are conducive for disease development
- Use in conjunction with protectant fungicides
- Always read and follow the label.

USE PATTERN

CROP	DISEASE	RATE	FOLIAR SPRAY VOLUME	APPLICATION TIMING
Apple trees, Pear trees, Crabapples	Powdery mildew (<i>Podosphaera leucotricha</i>)	9 L/ha	Use minimum spray volume to obtain a complete coverage of foliage (500-800 L/ha)	Apply when conditions are conducive for disease development
	Apple scab (<i>Venturia inaequalis</i> , <i>Venturia pirina</i>)			Apply after rain or a prolonged period of moisture, when conditions are conducive for disease development (post infection; after spores are release up to
Grapevine	Powdery mildew (<i>Erysiphe necator</i>)	18 L/ha	Use minimum spray volume to obtain a complete coverage of foliage (500-800 L/ha)	Apply when conditions are conducive for disease development
Crop group 9B (including squash, cucumber)	Powdery mildew (<i>Sphaerotheca fuliginea</i> <i>syn. Podosphaera xanthii</i> <i>Erysiphe cichoracearum</i>)	18 L/ha	Use minimum spray volume to obtain a complete coverage of foliage (700 L/ha)	Apply when conditions are conducive for disease development



The Use of Agral or Ag-Surf in tank mix with Mancozeb fungicide is not recommended.

Buran is registered in Canada by AEF Global. Please refer to product label for complete application details. Always read and follow label directions.

Log on to www.aefglobal.com for full label information.