











TECHNICAL SPECIFICATIONS

• Active ingredients: Citric acid

• Packaging: One (1) case = 2×2.5 gal

• Formulation Type: Liquid concentrate

• Chemical Group: N/A Biopesticide

• PHI: 0 days

• REI: When spray is dry or 4 hours

MODE OF ACTION

FungOUT forms a protective barrier on leaf surface that inhibits pathogen development. It also has antifungal effects on conidia germination, germ tube formation and elongation, appressorium formation, hyphae branching and haustorium formation.

In addition, it has a direct bactericide activity and also reduces bacterial growth and the appearance of various disease symptoms.

Manufactured by:

925 des Calfats • Lévis (QC) G6Y 9E8

Toll free: 1 (866) 622-3222 • Tel.: (418) 838-4441 • Fax: (418) 838-9909

Update: 26/09/2018 info@aefglobal.com • www.aefglobal.com





USE PATTERN

CROP	DISEASE	RATE	FOLIAR SPRAY VOLUME	APPLICATION TIMING
Greenhouse and field tomatoes	Bacterial canker (Clavibacter michi- ganensis)	2.1% v/v	Use minimum spray volume to obtain full coverage	Apply preventatively: Seedling emergence, before grafting, after grafting, before transplant shipping, at reception of transplants reception and once a month thereafter Apply at first sign of disease, 5 to 7 days interval, to contain the progression of the disease.
	Powdery mildew (Oïdium neolycoper- sici)	1.2 to 1.6% v/v		Apply at first sign of disease.

FungOUT has not been registered by the United States Environmental Protection Agency. AEF Global Inc. represents that this product qualifies for exemption from registration under the Federal Insecticide, Fungicide and Rodenticide Act. Please refer to product label for complete application details.

Always read and follow label directions. Log on to www.aefglobal.com for full label information.

AEF GLØBAI