

BERRY CARE

FUNGICIDE AND BACTERICIDE CONCENTRATE

FOR OUTDOOR FOOD AND GREENHOUSE USES



Active ingredient: Citric acid.....	1.07%
Inert ingredients: Water, lactose, ethyl lactate, potassium sorbate, sodium benzoate.....	98.93%
Total:	100.00%

NET CONTENTS:

2.5 Gallons

Batch:

This product has not been registered by the United States Environmental Protection Agency. AEF Global Inc. represents that this product qualifies for exemption from registration under the Federal Insecticide, Fungicide and Rodenticide Act.

CAUTION / KEEP OUT OF REACH OF CHILDREN / READ THE LABEL BEFORE USING

See back panel for additional Precautionary Statements

11/11/2020 v1



Manufactured by:

AEF GLOBAL Inc.

925, des Calfats, Lévis, QC, G6Y 9E8

Phone: (418) 838-4441

Toll free help-line: 1-866-622-3222

www.aefglobal.com

FIRST AID

If eye or skin contact occurs, immediately flush eye or rinse skin with plenty of water. If swallowed, rinse mouth with clean water. Call a Poison Control Center or doctor for treatment advice.

PRECAUTIONARY STATEMENTS

CAUTION. May cause eye and skin irritation. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear long-sleeved shirt and long pants, waterproof gloves, eye goggles and shoes plus socks. Keep and wash PPE separately from other laundry. Wash with detergent and hot water.

DIRECTIONS FOR USE

Shake well before use. Apply as a foliar treatment for crops and at the application rates listed below. For maximum effectiveness, apply BERRYCARE when conditions are conducive to disease development, or at the early stages of disease development. Use a combination of spray volume and pressure to obtain an excellent coverage of both sides of leaves. Avoid spraying when significant rainfall is expected within 24 hrs. The product can leave a white or shiny hydrosoluble residue on plants; the use of a silicone-based surfactant at the rate of 2 fl oz per 100 gallons greatly reduces the presence of residues. It is however recommended to evaluate the compatibility of each surfactant on a small group of representative plants before applying to a large number of plants. Do not mix with alkaline products. BERRYCARE contains weak acid and could react with products containing significant amounts of carbonates

Restricted Entry Interval: After a BERRYCARE application, you may reenter the area when the product is dry or after 4 hours. **Pre-Harvest Interval:** BERRYCARE can be applied up to and including the day of harvest.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Pesticide Storage: Keep at room temperature, away from direct sun. Keep from freezing. Store in original container out of reach of children. Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. Container Handling: Nonrefillable container: Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into the application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after flow begins to drip. Repeat this procedure two more times. Offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

WARRANTY

To the extent consistent with applicable law, seller makes no warranty, express or implied, of merchantability, fitness or otherwise concerning the use of this product other than as indicated on the label. User assumes all risks of use, storage or handling not in strict accordance with accompanying directions.

DILUTION TABLE PER ACRE

Rate	Spray volume		
	25 gpa ¹	50 gpa	100 gpa
1.4%	0.4 gal	0.7 gal	1.4 gal

¹ Gallons Per Acre

APPLICATION RATES

Crop	Disease	Rate	Interval of application
Crop group 13-07 Small Fruits and Berries such as: Blackberries, Currants, Grapes, Strawberries, Raspberries, Cranberries, Highbush and Lowbush Blueberries	Powdery mildew (<i>Podosphaera aphanis</i> syn <i>Sphaerotheca macularis</i> f.sp.fragariae, <i>Erysiphe necator</i>) Anthracnose (<i>Colletotrichum acutatum</i>)	1.4 % v/v with a non-ionic (spreader) surfactant	5 – 10 days
	Angular leaf spot (<i>Xanthomonas fragariae</i>)	1.4 % v/v	

